

occurs because routine TB slaughter surveillance has limited detection capabilities. The estimated sensitivity of slaughter surveillance for beef herds in the US ranges from 3-7%, and the probability of detecting an affected beef herd within five years ranges from 15-35%, depending on herd size (USDA, Animal and Plant Health Inspection Service, Veterinary Services, Center for Epidemiology and Animal Health 2009). Mexican origin feeder and rodeo cattle are listed as one of the three most likely sources of TB introduction in the US (Camacho, 2021).

Despite tremendous efforts to eradicate *Mycobacterium bovis* from the US cattle herd for nearly a century, novel strains of TB continue to emerge in western states in beef and dairy herds with inconclusive epidemiological investigations. With the availability of EIDs, the US has an opportunity to make important changes to identification requirements for Mexican rodeo cattle, align import rules with those of Canada, and improve traceability to safeguard the domestic cattle herd.

Resolution:

The United States Animal Health Association (USAHA) urges the United States Department of Agriculture (USDA), Animal and Plant Health Inspection Service, Veterinary Services to require electronic 484 prefix official identification ear tags in addition to "M" branding for all rodeo type cattle born in Mexico (regardless of end use) that enter the United States. To further improve animal disease traceability, USAHA urges USDA to provide guidance on how to officially identify Mexican origin cattle that lose ear tags.